

search of the Soviet and Eastern European literature on this subject alone, a total of about 100 publications authored by 75 researchers was found and this figure is probably a conservative reflection of the available works which are estimated to be several hundred.

In an attempt to summarize the prolific Soviet and Eastern European work on clinical and hygienic aspects of exposure to microwaves, it became apparent that a number of human systems and functions had been documented to be affected by this factor (Table 1). By far the most frequently and repeatedly reported human responses to microwaves involve the central nervous system. These responses have been noted for a wide range of frequencies ( $\sim 30$ – $300,000$  MHz) at both thermogenic ( $>10$  mW/cm<sup>2</sup>) and nonthermogenic (microwatts to milliwatts/cm<sup>2</sup>) intensities.

An often disappointing facet of the Soviet and East European literature on the subject of clinical manifestations of microwave exposure is the lack of pertinent data on the circumstances of irradiation; frequency, effective area of irradiation, orientation of the body with respect to the source, waveform (continuous or pulsed, modulation factors) exposure schedule and duration, natural shielding factors, and a whole plethora of important environmental factors (heat, humidity, light, etc.) In addition, the physiological and psychological status of human subjects such as health, previous or concomitant medication, and mental status is also more often than not omitted. These variables, both individually and combined, affect the human response to microwave radiation. Despite these omissions, however, the reviewer cannot help but be impressed both by the consistency of the findings and the large size of Soviet and East European clinical and hygienic surveys which have involved literally thousands of people over the past 20 or more years.

## CLINICAL MANIFESTATIONS OF EXPOSURE TO RADIOFREQUENCY FIELDS

### General Clinical Syndromes

Many Soviet clinical workers have attempted to categorize the chronological stages of human responses to microwaves. Panov et al. (12) proposed three categories or stages of responses to microwaves (Table 2). These were listed as the asthenic syndrome, characterized by fatigue, depression, and a number of other changes. This first stage is not

TABLE 1

*Effects of electromagnetic radiation on the human organism*

- |                                     |
|-------------------------------------|
| I. Central Nervous System           |
| II. Autonomic Nervous System        |
| III. Neurohumoral Systems           |
| IV. Endocrine Glands and Functions  |
| V. Eye and Ocular Functions         |
| VI. Blood and Hematopoietic Systems |
| VII. Miscellaneous Organs           |

marked by severe episodes such as fainting or dramatic changes in pulse or blood pressure and the subject responds to outpatient treatment. The second category is called the "syndrome of autonomic and vascular dystonia". The essential feature of this stage is pulse lability (brady- and tachycardia), blood pressure lability (hypo- or hypertension), EKG changes, and general neurocirculatory asthenia. Severe episodes such as fainting spells may occur and the subject requires hospitalization of unspecified nature or duration. The third stage is referred to as the diencephalic syndrome in which visceral dysfunctions and crises are observed. Typical episodes during this stage are listed as "apathic amblic" disorders, hypersomnia, hypokinesia, hypothalamo-pituitary-suprarenal weakness, and inhibition of sexual and digestive reflexes. Panov claims that these changes are not always reversible and that subjects require hospitalization. It should be noted that Panov did not specify the nature or duration of outpatient or hospital treatment, nor did he relate these symptoms to specific irradiation parameters.

### General Subjective Complaints (Indirect Effects on the CNS)

A large number of East European and especially Soviet clinical and hygienic workers (13–22) have consistently and repeatedly documented an astonishing number of subjective complaints which are usually referred to as evidence of the direct or indirect effect of microwaves on the central nervous system (Table 3). These responses have been reported for a wide range of wavelengths (30– $300,000$  MHz) and field intensities (microwatts to several milliwatts/cm<sup>2</sup>). Unfortunately, it is often difficult to attach any significance to Soviet clinical findings in the absence of pertinent data on exposures and on patient backgrounds. Typical, for instance, was a survey conducted by Sadchikova (21) in which three groups of occupational personnel (technicians, assemblers, and maintenance workers around centi-

TABLE 2

*Soviet classification of general clinical syndromes of exposure to electromagnetic radiation EMF's*

- |  |
|--|
| A. The Asthenic Syndrome (reversible; outpatient treatment)    |
| 1. fatigability and emotional changes                          |
| 2. acrocyanosis  |
| 3. increased perspiration of extremities                       |
| 4. increased pilomotor reflex                                  |
| 5. dermatographism   |
| 6. pulse lability  |
| 7. blood pressure lability                                     |
| B. Autonomic Cystonia (reversible; hospitalization)            |
| 1. hyper- or hypotension                                       |
| 2. bradycardia and tachycardia                                 |
| 3. changes in EKG signs  |
| 4. fainting spells   |
| C. Diencephalic Syndrome (usually reversible; hospitalization) |
| 1. insomnia  |
| 2. adynamia  |
| 3. hypothalamo-pituitary-suprarenal inhibition                 |
| 4. inhibition of sexual function and digestive reflexes        |

meter wave generators) were exposed to: (1) periodic intense radiation (3–4 mW/cm<sup>2</sup>); (2) moderate radiation (tenths of mW/cm<sup>2</sup>); and (3) weak radiation (hundredths-tenths of mW/cm<sup>2</sup>). As can be seen in Table 4, the group exposed to the weakest radiation was shown to display the highest incidence of complaints. This finding and lack of pertinent exposure data such as duration and affected body area make these data difficult to accept on face value. On the other hand, Edelwejn (14) has conducted interesting and comprehensive neurological examinations and interviews of Polish personnel exposed for up to six hours/day to microwave field intensities of 10 microwatts to several milliwatts/cm<sup>2</sup>. He found that many of the subjective complaints listed in Table 3 (headaches, dizzy spells, fatigue, perspiration, etc.) depended upon the length of employment and degree of exposure. Only subjects exposed to high (mW/cm<sup>2</sup>) intensities exhibited EEG changes. Edelwejn was of the opinion that there is a dramatic response to microwave exposure occurring during the first three years which are accompanied by neurotic symptoms. This three year period is followed by a phase of gradual adaptation. The reappearance of neurologic symptoms occurs after a long period (many years) of exposure to microwaves, even after adaptation has occurred.

Osipov (1965) (20) in a review of neurologic responses to microwave exposure concluded that most subjective symptoms were reversible and that patho-

logical damage to neural structures was insignificant. Only rarely were microwaves found to cause hallucinations, syncope, adynamia and other manifestations of the so-called "diencephalic" syndrome.

Soviet workers have also documented subjective complaints identical to those in Table 3 as a result of exposure to electric and magnetic fields. Vyalov et al. (23) reported characteristic microwave symptoms such as headache, fatigue etc., in workers exposed to 150–1500 oersted magnetic fields. Asanova (24) reported analogous findings for workers exposed to 115–125 microampere fields around hydroelectric stations.

#### *Functional Changes in the CNS*

Many Soviet and Eastern European workers have attempted to identify specific CNS functional responses to microwave exposure. Most Soviet workers are of the opinion that the CNS is the most sensitive of all systems to the effects of microwaves, both at thermogenic and nonthermogenic field intensities. Based primarily upon experimental research, Presman (9) is of the opinion that the hypothalamus is the most sensitive CNS structure to microwave effects which would explain, in his view, the high incidence of blood and humoral changes noted in human subjects exposed to this factor.

Changes in human CNS function have been evaluated on the basis of EEG surveys, reflex tests, and general neurological examinations (Table 5). These changes are reported for a wide range of frequencies and field intensities (thermal and nonthermal). However, functional CNS responses appear to be de-

TABLE 3

*General subjective complaints resulting from exposure to electromagnetic radiation*

- |   |
|---|
| 1. Pain in head and eyes                                      |
| 2. Lacrimation  |
| 3. Weakness, weariness and dizziness                          |
| 4. Depression, antisocial tendencies, general irritability    |
| 5. Hypochondria, sense of fear, and general tension           |
| 6. Impairment of memory and general mental function           |
| 7. Adynamia and inability to make decisions                   |
| 8. Inhibition of sex life (male)                              |
| 9. Scalp sensations and loss of hair                          |
| 10. Chest pain and heart palpitation                          |
| 11. Dyspepsia, epigastric pain, and loss of appetite          |
| 12. Trembling of eyelids, tongue, and fingers                 |
| 13. Asthma  |
| 14. Brittle fingernails                                       |
| 15. Sensitivity of mechanical stimulation and dermatographism |

TABLE 4

*Changes in the nervous system as a result of exposure to microwaves*

Group	No. examined	Changes observed (in % of subjects studied)							
		Headache	Increased fatigue	Increased irritability	Sleepiness	Delayed dermographism	Slowed orthostatic reflex	Wrist hyperdermosis	Thyroid hypertrophy
1	184	12	20	8	2	16	19	6	15
2	129	39	31	12	14	7	21	37	—
3	78	36	31	15	19	14	11	26	52
Control	100	8	10	8	2	—	—	4	14

pendent upon wavelength; direct effects on the brain were reported by Gordon (1964) (25) and Presman (9) to intensify with increase in wavelength. However, when reactions are due to a combination of peripheral and direct stimulation, it is impossible to correlate response with wavelength.

A number of workers have reported changes in EEG patterns as a result of exposure to microwaves. Klimkova-Deutschova (26), a Czechoslovakian researcher, reported that both clinical and EEG findings suggested a predominance of an inhibition process. EEG's showed a predominance of sleep rhythms. In this connection, the interesting (if rather curious) work of Ivanov-Muromskiy (27), a Soviet expert on electrosleep and electroanesthesia

deserves comment. His research on human subjects suggested that pulsed (10-1000 Hz) UHF fields of nonthermal intensity directed from bitemporal electrodes a few inches from the subject's head could induce inhibition similar to that produced by pulsed electrical currents (electrosleep). Unfortunately, this research was not described in detail by Ivanov-Muromskiy.

Drogichina (13) reported that CNS damage is characterized by the "asthenic syndrome" which can be detected from EEG and neurological findings. Presman (9), in reviewing Soviet, Czechoslovakian, and Polish work, reports that the EEG's of subjects exposed to weak (nonthermal) microwave field intensities show an increased incidence of slow, high amplitude waves. In Poland, Edelwejn and Baranski (14) reported a decreased incidence of alpha rhythms and a decreased percentage of alpha waves in subjects exposed to "high" ( $\text{mW}/\text{cm}^2$ ) intensities of microwave fields. All subjects examined in this study over-reacted to the administration of cardiozol, a respiratory and cardiac stimulant. In general, because of the rather primitive state-of-the-art of EEG analysis, these findings should be viewed with extreme caution.

Perceptual changes as a result of exposure to microwaves have also been frequently reported. Livshits (28) reported that "high intensity" microwaves had been found by Soviet workers to cause hallucinations. He also reported that high frequency, high intensity fields had been demonstrated to cause involuntary motor reactions in one healthy individual. Matuzov (29) noted visual perception changes after a 10 minute exposure to 10 cm microwaves of nonthermal ( $1.1 \text{ mW}/\text{cm}^2$ ) intensity. He found a considerable decrease in blind spot area,

TABLE 5

*Functional CNS changes resulting from exposure to electromagnetic radiation*

1. Changes in EEG patterns
  - a. "asthenic" signs
  - b. predominance of inhibition process
  - c. increased incidence of slow, high amplitude waves
  - d. decreased incidence of alpha rhythms and waves
  - e. predominance of "sleep" rhythms
2. Perceptual changes
  - a. hallucinations (visual)
  - b. decrease in ocular blind spot area
  - c. shortening of optic chronaxie and reduction of rheobase
  - d. auditory sensitivity changes
  - e. decreased olfactory sensitivity
  - f. increased olfactory activity
  - g. parapsychologic phenomena
3. Alternating arousal and drowsiness
4. Stimulation of motor functions
5. Depression of mental functions
6. Involuntary motor reactions

TABLE 6

*Autonomic and cardiovascular effects of electromagnetic radiation.*

1. Changes in cardiac function (EKG)
  - a. decreased spike amplitude
  - b. lengthened QRS interval
  - c. slowed auricular and ventricular conductivity
2. Bradycardia and tachycardia
3. Hyper- and hypotension
4. Increased precapillary resistance
5. Increased vascular elasticity

shortening of optic chronaxie, and reduction of rheobase in two subjects. These effects were judged to be nonthermal (specific) and were found to be reversible. Sheyvekhman (30) noted changes in auditory sensitivity (5–10 dB) in response to 6 meter waves pulse modulated at 300, 1000, or 4000 Hz applied for five minutes to the heads of human subjects. He did not clarify whether sensitivity was increased or decreased. Lobanova and Gordon (31) noted a decrease in olfactory sensitivity after exposure to microwaves and suggested that this response might be a good index for identifying harmful microwave effects. These authors also found an increase in olfactory excitability (decreased threshold) after a single dose of caffeine. This was suggested as evidence of functional olfactory changes caused by microwaves.

In the realm of parapsychology, it is interesting to note that leading Soviet researchers who strongly believe in the nonthermal CNS effects of microwaves are involved in the electromagnetic (centimeter wave) theory of extrasensory perception (3). This work, initiated in 1966, is being conducted for a special Bioinformation Section of the Scientific and Technical Society of Radiotechnology in Moscow. The results of Soviet ESP research have thus far been interesting but statistically inconclusive.

Both the stimulatory and inhibitory effects of microwaves on CNS function have been frequently documented by Soviet workers. Subbota (32) reported alternating arousal and drowsiness in response to microwaves in working with dogs. As mentioned earlier, the Soviet electrosleep expert, Ivanov-Muromskiy (27) concluded from his studies of human subjects that pulsed UHF fields could be used as a form of contactless electrosleep which he calls "radio-sleep". Depression of mental function, inability to concentrate, and general sluggishness is frequently documented by Soviet and Eastern European re-

searchers as a subjective response to microwave exposure.

#### *Autonomic and Cardiovascular Responses*

Reports of human autonomic and cardiovascular responses to microwaves are nearly as numerous as those documenting CNS responses to this factor (Table 6). Responses are noted for a wide range of frequencies at thermal and nonthermal field intensities and during acute and chronic exposure. Decreased EKG spike amplitudes have been noted by Drogichina (33) in subjects working around radio-frequency fields. Sadchikova (34) reported on various cardiovascular shifts in workers exposed to different field intensities (Table 7). Figar (15) and Smurova (35) have noted decreased coronary conductivity, sinusoidal arrhythmia, brady- and tachycardia, and oscillating hypo- and hypertension. Monayenkova et al. (36) studied minute blood volume, peripheral resistance, average arterial pressure, and smooth muscle tonus using a mechanocardiograph. She found that a tendency toward hypertension, increased elasticity of myogenous vessels, increased precapillary resistance, sinus bradycardia, and changes in intracardiac conductivity were more often noted in exposed than in unexposed subjects. All of these changes were found to be reversible with one or two questionable exceptions.

There is some evidence that certain enzymes implicated in CNS function might be affected by exposure to microwaves (Table 8). Revuts'kyy et al. (37) found a change in the specific cholinesterase activity of erythrocytes in human whole blood with 13.56 and 23.75 MHz microwaves. The 13.56 MHz radiation was found to decrease blood histamine content while not altering cholinesterase activity. The 23.75 MHz radiation did not change blood histamine content but increased cholinesterase activity. Bartonicek et al. (38) surveyed the blood biochemistry of workers exposed to centimeter waves. Of a total of 27 blood sugar curves, 7 were flat, 7 were prediabetic, and four indicated slight glycosuria. The distribution of pyruvic and lactic acid and creatinine are shown in Table 9. Lactic acid was found to be decreased 2.5 times more than it was found to be increased. Roughly 75% of the subjects exposed to microwaves and examined by Bartonicek were reported to have prediabetic blood sugar curves. These metabolic shifts were attributed to autonomic dysregulation, possibly indicative to diencephalic lesions resulting from early exposure to centimeter

TABLE 7

*Cardiovascular changes in subjects exposed to electromagnetic radiation (Sadchikova, 1964)*

Range	EMF parameters	Exposure/control ratio		
	Field intensity	hypertonia	bradycardia	increase of QRS interval (up to 0.1 sec)
SHF	1-several mW/cm <sup>2</sup>	1.85	24.0	11.5
	1 mW/cm <sup>2</sup>	2.0	16.0	12.5
UHF LF	nonthermal	1.2	8.0	21.0
	tens to hundreds V/M	9.21	12.0	—
	hundreds to 1000 V/M	1.2	5.0	—
Percent incidence in controls		14%	3%	2%

waves. Gel'fon and Sadchikova (39) noted increased blood globulins in 50% of a group exposed to microwaves which indicated a shift in the albuminglobulin coefficient. Haski (40) noted slight changes in the levels of blood sugar, cholesterol, and lipids of healthy subjects exposed to microwaves. However, there was a pronounced decrease in all three categories when diabetics were exposed.

#### *Hematopoietic and Biochemical Responses*

Numerous human hematopoietic changes have been reported to result from exposure to microwave fields (Table 10). The severity of these changes range from minimal to significant. Sokolov (41) noted reticulocytosis in radar workers. Baranski (42) observed that a small drop in erythrocytes occurs in all people exposed to microwaves and that the phenomenon is related to the duration and severity of exposure. About 50% of the subjects examined by Baranski showed a moderate decrease in

platelet count. Lysina (43) noted basophilic granularity of erythrocytes and was of the opinion that this index should be taken as an initial sign of microwave effects on the human organism. Presman et al. (44) found that the osmotic resistance of erythrocytes was negatively affected by microwaves. Smurova (22) and others found that the shape and volume of erythrocytes changes as a result of exposure to microwave fields. Prolonged exposure was occasionally noted to result in hemolytic processes. An increase in the RNA level of lymphocytes was also noted by Smurova in workers chronically exposed to microwaves; this finding corresponded to a concomitant increase in monocytes (young cells) which contain the greatest quantity RNA. Baranski (42) detected various leukocyte shifts in workers exposed for one year to microwaves. Normalization of this index was found to occur after prolonged exposure to this factor. He also found a tendency towards lymphocytosis with accompanying eosinophilia in subjects exposed for more than five years to low and moderate microwave intensities.

Soviet workers have also found biochemical changes to occur in other sites (8). A drop in RNA content was noted in the spleen, liver, and brain in animals chronically exposed to microwaves while DNA content was found to remain constant.

#### *Ocular Responses*

Changes in human ocular function and eye pathology are widely documented and occur primarily

TABLE 8

*Neurohumoral responses to radiofrequency electromagnetic radiation*

1. Altered cholinesterase activity in human whole blood (erythrocytes)
2. Decrease in blood histamine content
3. Increase in blood proteins
4. Altered carbohydrate metabolism
5. Changes in blood sugar, cholesterol, and lipids (pronounced in diabetics)
6. Decreased hemoglobin

TABLE 9

*Distribution of pyruvic and lactic acid and creatinine excretions in workers exposed to microwaves*

	Pyruvic acid		Lactic acid		Creatinine	
	number	%	number	%	number	%
No. of measurements	40	100.0	35	100.0	34	100.0
Normal	28	70.0	14	40.0	14	41.2
Increased	4	10.0	6	17.2	6	17.6
Lowered	8	20.0	15	42.8	14	41.2
Averages	0.77 mg%		14 mg%		1.33 mg%	
Controls	0.65 mg%		17 mg%		1.30 mg%	
Established standard	0.5—1.0		10—20		1.2—1.9	

after acute or chronic exposure to thermogenic microwave intensities (Table 11). As mentioned earlier, one Soviet worker (28) has reported that exposure to intense microwave fields was noted to cause hallucinations. Matuzov (29) found the area of the blind spot to decrease after exposure to nonthermogenic (10 cm; 1.1 mW/cm<sup>2</sup>) microwave field intensities. Other Soviet workers, as reported by Marha (8), have found that microwave radiation (a few mW/cm<sup>2</sup>) can cause a decrease in sensitivity to color (blue) and difficulty in detecting white objects. Changes in intraocular pressure have also been noted by Soviet workers as have altered sensitivity to light stimuli during exposure to pulsed and nonpulsed fields. General ocular pain, eye strain and fatigue, eyelid tremor, and lacrimation are also common symptoms noted by Soviet workers.

Pathological changes in the eye (cataracts) occur primarily as a result of exposure to thermogenic (greater than 10 mW/cm<sup>2</sup>) microwave intensities. Sadchikova (45) and other Soviet workers (6) have noted unilateral and bilateral cataracts to occur in subjects exposed to several mW/cm<sup>2</sup> field intensities. Presman (44) noted a drop in vitamin C content in the lens and anterior chamber fluid at nonthermogenic intensities. In the event of acute cataract development a decrease in ATP and pyrophosphatase activity of the lens was noted. In addition, it is suspected that damage to tissue respiration and oxida-

tion mechanisms as a result of exposure to microwaves can lead to cataract formation.

There is some evidence that ocular responses to microwaves are frequency dependent. Pol (46) noted that 10 GHz fields caused anterior lens opacity while 2.45 GHz cause posterior opacity.

Belova (47) noted that in 370 microwave generator workers exposed to mW/cm<sup>2</sup>, lacrimation, ocular fatigue, and frequent conjunctival irritation would occur at the end of each working day. Zydecki (48) suggested that all candidates for occupation around microwave sources receive comprehensive ophthalmological examinations. This suggests that certain ophthalmological profiles might be more vulnerable to microwave radiation than others.

TABLE 10

*Hematopoietic and biochemical responses to electromagnetic radiation*

1. Blood
  - a. reticulocytosis
  - b. basophilic granularity of erythrocytes
  - c. decrease in erythrocytes, platelets and hemoglobin
  - d. altered osmotic resistance of erythrocytes
  - e. neutrophilic leukocytosis
  - f. lymphocytosis, monocytosis, and eosinophilia
  - g. increased RNA in lymphocytes
2. Organs
  - a. Decreased RNA content in brain, liver, and spleen

*Endocrine Responses*

Damage to sex glands and functions have frequently been documented to occur after chronic exposure to primarily thermal microwave fields (Table 12). Marha (8) in reviewing Soviet and East European findings noted decreased spermatogenesis, altered sex ratio of births, changes in menstruation, retarded fetal development, congenital effects in newborn babies, decreased lactation in nursing mothers, and other related responses to occur as a result of exposure to thermal (i.e., greater than 10 mW/cm<sup>2</sup>) microwave intensities. Microwaves were also implicated in an increase in the percentage of miscarriages in both humans and animals. Some of these

TABLE 11

*Effects of electromagnetic radiation on the eye*

1. Perceptual and function changes
  - a. hallucinations
  - b. decrease in size of blind spot
  - c. decreased sensitivity to color (blue)
  - d. difficulty in detection of white objects
  - e. decreased sensitivity to light stimuli in dark adapted eye
  - f. change in intraocular pressure
  - g. lacrimation, ocular fatigue, and ocular pain
  - h. trembling of the eyelids
2. Pathological changes
  - a. lens coagulation (cataracts)
  - b. decrease in vitamin C content of lens and vitreous humor
  - c. decrease in ATP and pyrophosphinase activity
  - d. anterior and posterior lens opacity
  - e. conjunctival irritation

findings reported by Marha are consistent with subjective complaints reported by Soviet researchers such as decreased sex activity, mentioned earlier. Specific genetic changes resulting from exposure to either thermal or nonthermal microwave fields have yet to be demonstrated.

Soviet sources have reported pituitary and other endocrine responses to microwave exposure. Kolesnik (49) suggested that pituitary-hypophyseal-adrenal changes were primarily due to CNS influences on the hypophysis after exposure to microwaves. Drogichina (33, 50), Sadchikova (21, 34), and Smirnova (51) have reported thyroid gland enlargement and increased iodine-131 uptake. These changes suggest an increase in thyroid stimulating hormone (6). Hasik (40) and Presman (44) noted increased activity of the adrenal cortex to occur after microwave ex-

TABLE 12

*Endocrine responses to radiofrequency radiations*

1. Sex organs and ontogenesis
  - a. thermal trauma (tissue damage) to male reproductive tissues
  - b. decreased spermatogenesis (sterility)
  - c. altered sex ratio of births (more girls)
  - d. altered menstrual activity
  - e. altered fetal development
  - f. decreased lactation in nursing mothers
2. Endocrine glands
  - a. altered pituitary and pituitary-hypophyseal function (CNS)
  - b. hyperthyroidism
  - c. thyroid enlargement
  - d. increased iodine-131 uptake
  - e. increased adrenal cortex activity
  - f. decreased corticosteroids in blood
  - g. decreased glucocorticoidal activity

posure. Murashov (52) studied 20 subjects occupationally exposed to UHF fields. He noted a reduction in plasma corticosteroid content which was attributed to lowered adrenal, or possibly sex gland androgenic activity.

*Miscellaneous Responses*

Loshak (53) reported that various human responses, such as subjective complaints as a result of chronic microwave exposure, appeared to vary slightly with climate (Table 13). In general, responses to microwave fields were more pronounced in hot, dry climates. It was found that the electrical resistance of the skin of exposed workers was lower than in unexposed workers in a hot climate. Decreased resistance was attributed both to CNS stimulation or increased sympathetic tonus due to skin receptor reactions. These findings, while not dramatic, led Loshak to speculate that special hygienic considerations for workers exposed to microwaves in a hot climate should be exercised (improved ventilation etc.).

TABLE 13

*Miscellaneous effects on electromagnetic radiation*

1. Climatic effects
  - a. responses to electromagnetic radiation more pronounced in hot climate
  - b. decreased electrical resistance of skin in hot climate due to electromagnetic radiation
2. Internal Organs
  - a. dyspepsia and epigastric pain
  - b. decreased appetite
  - c. liver enlargement

Orliva (54) and others have reported that workers exposed to microwaves complain of decreased appetite, dyspepsia, pain in the epigastric region, and exhibit enlargement of the liver. Marha (8) in reviewing Soviet and Czechoslovakian experimental work on animals reported that exposure to microwaves was noted to cause liver hemorrhaging, hepatic cell degeneration, and decreased filtration or renal tubules. Analogous findings for humans have not been documented.

### CONCLUSIONS

The large body of the Soviet and East European clinical and hygienic findings on human responses to microwave radiation reviewed in this paper suggest that a surprisingly wide variety of neurological and physiological reactions are to be expected during exposure to nonthermal (i.e., less than 10 mW/cm<sup>2</sup>) field intensities within an extremely wide range of frequencies (approximately 30–300,000 MHz). These reactions, which are generally reversible, are often documented as a result of human exposure to field intensities as low as a few microwatts/cm<sup>2</sup>. They are reported to be primarily effects upon the nervous system and reflect traditionally heavy Soviet emphasis on the central nervous system. Soviet and East European findings in this area are therefore in striking contrast to those of the West which have, in the main, documented non-CNS responses to thermal (i.e., greater than 10 mW/cm<sup>2</sup>) intensities. Only in the realm of human endocrine, visual, and skin receptor responses to thermal microwave burdens is any real substantive agreement between Soviet and Western findings to be found.

The substantially lower Soviet and East European daily maximum permissible dose (MPD) value for human exposure to microwave radiation (0.01 mW/cm<sup>2</sup> vs 10 mW/cm<sup>2</sup> in the U.S.) appears to be based upon extensive findings of human subjective and other CNS-related responses to extremely low microwave field intensities and upon considerable CNS-oriented research on animals conducted in those countries. These findings also indicate that extensive dosimetric surveys around industrial and military sources of microwave radiation have been conducted in those countries (see, e.g., the report of Marha in this collection), although the extent and nature of Soviet work in this specific area has not been well documented. In the general context of differing U.S. and Soviet MPD's and MPC's, it is

important to note that the Soviet union has traditionally been more conservative with regard to many industrial hazards than the U.S.

Although the majority of Soviet findings on human responses to low intensity microwave fields must be regarded with extreme caution because of the omission of exposure and other pertinent data, it is suggested that the surprising consistency of this large body of findings merits the critical attention of the U.S. scientific community. Of particular interest is the relatively recent Eastern European work in this area. Research conducted in these countries, although heavily influenced by the Soviet Union in the early stages, appears to be of high quality; reflects a good awareness of both Soviet and Western approaches to the problem of the biological effects of microwaves; and suggests a trend towards more independent approaches to this problem. The fact that East European countries such as Czechoslovakia and Poland have adopted essentially the same maximum daily permissible dose for human exposure to microwaves as the Soviet Union is of interest and should be investigated in more detail. The Czechoslovakian MPD for microwaves, while admittedly (by Marha) influenced by the Soviet MPD, was arrived at only after considerable hygienic and dosimetric survey work had been conducted in that country. Nonetheless, it is suggested that until additional research on this difficult problem has been conducted in this country, and a more critical analysis of the available Soviet and East European findings has been made, a judgment of the U.S. 10 mW/cm<sup>2</sup> MPD is presently rendered difficult, if not impossible.

### REFERENCES

1. C. H. Dodge: ATD Report P-65-17, Library of Congress, 1965, 44 pp.
2. C. H. Dodge: ATD Report P-65-68, Library of Congress, 1965, 93 pp.
3. C. H. Dodge and S. Kassel: ATD Report 66-133, Library of Congress, 1966, 33 pp.
4. C. H. Dodge: Foreign Sci. Bull. 1, 7 (1965) (Library of Congress).
5. W. Thompson, et al.: ARL-TR-65-20, November 1965, 60 pp.
6. S. Michaelson, et al.: RAD-TR-67-461, September 1967, 63 pp.
7. J. Healer, et al.: ARA-348-1, April, 1967, 11 pp.
8. K. Marha, et al.: Prague, Statni zdravotnicki nakladatelstvi, 1968, 304 pp.
9. A. Presman: *Electromagnetic Fields and Life*, Moscow, Izd-vo Nauka, 1968, 288 pp.
10. S. Turlygin: Dokl. Akad. Nauk (SSSR), Nov. Ser. 17 (1-2), 11 (1937).



20. N. N. Livshits: *Biofiz.* No. 3, p. 378 (1957).
21. A. G. Panov, et al.: *Voyenno-med.* No. 9, p. 13 (1966).
22. E. A. Drogichina, et al.: *Tr. Profzabol.* No. 1, p. 17 (1965).
23. A. Edelwejn, et al.: *Lek. Wojskowy*, No. 9, p. 781 (1966).
24. S. Figar: *Cesk. Physiol.* No. 5, p. 316 (1963).
25. Z. V. Gordon: *Vest. Akad. Med. Nauk (SSSR)* No. 4, p. 22 (1960).
26. V. Gur'yev: *Fitness, Expertise, and Work Arrangements for Nervous and Mental Diseases*, Vol. 18, p. 121 (1965) (JPRS 36, 164).
27. S. Haduch, et al.: *Lek. Wojskowy* 36, 119 (1960).
28. F. E. Kolesnik, et al.: *Voyenno-med. Zh.*, No. 4, 1967, p. 21 (1967).
29. Ya. A. Osipov: *Occupational Hygiene and the Effects of Radiofrequency Electromagnetic Fields on Workers*, Izd-vo medicina, Leningrad, 1965, p. 104.
30. M. N. Sadchikova: *The Biological Action of Ultrahigh Frequencies*, Izd-vo AN SSR, Moscow, 1960, p. 32.
31. Ye. I. Smurova: *Gig. Tr. Profzabol.* No. 5, p. 22 (1962).
32. A. M. Vyalov, et al.: *Gig. Tr. Profzabol.* No. 5, p. 39 (1966).
33. T. P. Asanova, et al.: *Gig. Tr. Profzabol.* No. 5 (1966).
34. Z. V. Gordon: *Biological Effects of Radiofrequency Electromagnetic Fields*, Izd-vo AMN SSSR, No. 2, Moscow, p. 3 (1964).
35. E. Klimkova-Deutschova: *Gewerbehyg* 20, 1 (1963).
36. K. A. Ivanov-Muromskiy: *Electrical Anesthesia and Electro-sleep in Man and Animals*, Naukova Dumka, Kiev, 1966, 221 pp.
37. N. N. Livshits, *Siofizika* No. 4, p. 426 (1968).
38. N. I. Matuzov: *Bull. Eksper. Biol. Med.* No. 7, p. 27 (1959).
39. B. E. Sheyvekhman, *Probl. Fiziol. Akus.* 1, 122 (1949).
40. Ye. A. Lobanova, et al.: *The Biological Action of Ultrahigh Frequencies*, Izd-vo AN SSR, Moscow, 1960, p. 52.
41. A. G. Subbota: *Bull. Eksper. Biol. Med.* No. 10, p. 55 (1958).
42. E. A. Drogichina: *The Biological Action of Ultrahigh Frequencies*, Izd-vo AN SSR, Moscow, 1960, p. 29.
43. M. N. Sadchikova: *Biological Effects of Radiofrequency Electromagnetic Fields*, Izd-vo AMN SSSR, Moscow, 1964, p. 110.
44. Ye. I. Smurova, et al.: *Gig. Sanitari.* No. 12, p. 27 (1964).
45. A. M. Monayenkova, et al.: *Gig. Tr. Profzabol.* No. 7, p. 18 (1966).
46. Ye. L. Revuts'kyy: *Fiziol. Zh. Akad. Nauk Ukr. SSR* No. 10, p. 379 (1964).
47. V. Bartonicek, et al.: *Casopis Lekaru Ceskych* No. 1, p. 26 (1964).
48. I. A. Gel'fon, et al.: *Biological Effects of Radiofrequency Electromagnetic Fields*, Izd-vo AMN SSSR, Moscow, 1964, p. 133.
49. J. Haski, et al.: *Pol. Tyg. Lek.* 15, 817 (1960).
50. V. V. Sokolov, et al.: *The Biological Action of Ultrahigh Frequencies*, Izd-vo AMN SSSR, Moscow, 1960, p. 43.
51. S. Baranski, et al.: *Lek. Wojskowy* No. 10, p. 903 (1966).
52. G. G. Lysina: *Gig. Sanitar.* No. 6, p. 95 (1965).
53. A. S. Presman, et al.: *Usp. Sovrem. Biol.* 51, 82 (1961).
54. M. N. Sadchikova, et al.: *Gig. Tr. Profzabol.* 6, 16 (1958).
55. W. Pol: *Lek. Wojskowy* 38, 318 (1962).
56. S. F. Belova: *Nauchn. Issled. Inst. Gig. Tr. Profzabol.* Tr. No. 1, p. 36 (1960).
57. S. Zydecki: *Lek. Wojskowy* No. 2, 124 (1967).
58. F. A. Kolesnik, et al.: *Voyenno-med. Zh.*, No. 4, 21 (1967).
59. E. A. Drogichina, et al.: *Gig. Tr. Profzabol.* No. 1, p. 28 (1962).
60. M. I. Smirnova, et al.: *The Biological Action of Ultrahigh Frequencies*, Izd-vo AMN SSSR, Moscow, 1960, p. 50.
61. B. F. Murashov: *Voyenno-med. Zh.* No. 6, p. 82 (1966).
62. A. Ya. Loshak: *Gig. Sanitar.* No. 6, p. 18 (1965).
63. A. A. Orlova: *The Biological Action of Ultrahigh Frequencies*, Izd-vo AMN SSSR, Moscow, 1960, p. 36.

BIOLOGICAL AND HEALTH EFFECTS OF ELECTROMAGNETIC (NONIONIZING) RADIATION  
Compiled by Hisako Halasz

TB 86-5

November 1986

**SCOPE:** The environment we live in today is filled with man-made electromagnetic fields generated by a variety of sources, including radio and TV transmitters and office visual display terminals. In addition to these artificially created electromagnetic fields, there also exists a natural background of electromagnetic fields.

Electromagnetic energy may vary greatly in frequency, i.e., wave length ranging from low energy, ultra-low frequency to high energy cosmic rays. The higher energy portion of the electro-magnetic spectrum is generally referred to as "ionizing radiation", and includes cosmic, gamma and x-rays. The lower portion of the spectrum is referred to as "nonionizing radiation" and includes ultraviolet (slightly ionizing at the higher end), visible and infrared light, microwaves, radiowaves and lower frequencies. Because of its low energy levels, concern over the health effects of nonionizing radiation is not as widespread as it is for ionizing radiation.

The term "electromagnetic pollution" is often used to describe the recent proliferation of artificial sources of electromagnetic fields in our environment. Sources referred to by this term generally are exclusive of ultraviolet, visible and infrared light. The electromagnetic spectrum covered by this guide follows the narrower definition above, namely, nonionizing electromagnetic fields from extremely low frequencies (ELF) to just below infrared.

While an extensive body of scientific and epidemiologic data has been accumulated, adverse biological and health effects of electromagnetic fields, thus defined, have not conclusively been demonstrated, with the exception of microwave thermal effects at higher intensities.

The following guide complements two previous guides, Low-Level Ionizing Radiation--Health Effects (TB 80-5) and Microwaves (TB 79-6) and is intended to help the interested reader locate materials on the biological and health effects of electro-magnetic fields in the collections of the Library of Congress. Not meant to be a comprehensive bibliography, it is designed--as the name of the series implies--to put the reader "on target."

## INTRODUCTIONS

Lerner, Eric J. Biological effects of electromagnetic fields. IEEE spectrum, v. 21, May 1984: 57-69. TK1.I15 and Pamphlet box\*

Morgan, M. Granger, and others. Power-line fields and human health. IEEE spectrum, v. 22, Feb. 1985: 62-68. TK1.I15 and Pamphlet box\*

SUBJECT HEADINGS used by the Library of Congress, under which books on the biological and health effects of electromagnetic radiation can be located in most card, book, and online catalogs, include the following:

ELECTRIC FIELDS--PHYSIOLOGICAL EFFECT (Highly relevant)  
 ELECTRIC LINES--OVERHEAD--HYGIENIC ASPECTS (Highly relevant)  
 ELECTRIC POWER DISTRIBUTION--HIGH TENSION--HYGIENIC ASPECTS (Highly relevant)  
 ELECTRIC WELDING--HYGIENIC ASPECTS (Highly relevant)  
 ELECTROMAGNETISM--HYGIENIC ASPECTS (Highly relevant)  
 ELECTROMAGNETISM--PHYSIOLOGICAL EFFECT (Highly relevant)  
 MAGNETIC FIELDS--PHYSIOLOGICAL EFFECT (Highly relevant)  
 MICROWAVES--HYGIENIC ASPECTS (Highly relevant)  
 MICROWAVES--PHYSIOLOGICAL EFFECT (Highly relevant)  
 RADIO WAVES--HYGIENIC ASPECTS (Highly relevant)  
 RADIO WAVES--PHYSIOLOGICAL EFFECT (Highly relevant)  
 RADIATION--PHYSIOLOGICAL EFFECT (Relevant)  
 RADIATION, BACKGROUND--PHYSIOLOGICAL EFFECT (Relevant)

See also names of individual instrument or devices which are capable of emitting electromagnetic radiation, when followed by subject subdivision "HYGIENIC ASPECTS" or "PHYSIOLOGICAL EFFECT," e.g.,  
 VIDEO DISPLAY TERMINALS--HYGIENIC ASPECTS

## GENERAL TEXTS

Becker, Robert O., and Andrew A. Marino. Electromagnetism and life. Albany, N.Y., State University of New York Press, c1982. 211 p.  
 Includes bibliographical references. QP82.2.E43B42\*

Brodeur, Paul. The zapping of America: microwaves, their deadly risk, and the coverup. New York, Norton, c1977. RA569.3.B76  
 Includes bibliographical references.  
 "Part of this book appeared originally as a series of articles in the New Yorker."

DeMatteo, Bob. Terminal shock: the health hazards of video display terminals. Toronto, NC Press; Port Washington, N.Y., distributed in the U.S.A. by Independent Publishers Group, 1985. 224 p.  
 RC965.V53D46 1985

Steneck, Nicholas H. The microwave debate. Cambridge, Mass., MIT Press, c1984. 279 p. RA569.3.S74 1984  
 Bibliography: p. 244-270.

\*Available in reference collection, Science Reading Room

ADDITIONAL TITLES

American national standard safety levels with respect to human exposure to radio frequency electromagnetic fields, 300 kHz to 100 GHz. Co-secretariat, Institute of Electrical and Electronics Engineers and U.S. Department of the Navy (Naval Electronic Systems Command). New York, the Institute of Electrical and Electronics Engineers, c1982. 24 p. ANSI C95.1-1982\*\*  
 Approved July 30, 1982, American National Standards Institute.

Biologic effects of environmental electromagnetism. Herbert L. Konig ... et al.. Translation by Terry C. Telger. New York, Springer-Verlag, c1981. 332 p. QP82.2.E43B54  
 Bibliography: p. 296-323.

Biological effects of electromagnetic radiation. Edited by John M. Osepchuk; associate editors, James W. Frazer ... et al.. New York, IEEE Press; sole worldwide distributor (exclusive of the IEEE), Wiley, c1983. 593 p. QP82.2.E43B554 1983\*  
 "Prepared under the sponsorship of the IEEE Committee on Man and Radiation."  
 Includes bibliographies.

Extremely low frequency (ELF) fields. Published under the joint sponsorship of the United Nations Environment Programme, the World Health Organization, and the International Radiation Protection Association. Geneva, W.H.O.; Albany, N.Y., distributed by WHO Publications Centre USA, 1984. 131 p. (Environmental health criteria, 35) RA569.3.E98 1984  
 Includes bibliographies.

Michaelson, Sol M. Microwave and radiofrequency radiation. Copenhagen, Regional Office for Europe, World Health Organization, 1977. 98 p. QP82.2.M5M5  
 "ICP/CEP 803."  
 Bibliography: p. 70-96.

National Council on Radiation Protection and Measurements. Biological effects and exposure criteria for radiofrequency electromagnetic fields: recommendations of the National Council on Radiation Protection and Measurements. Bethesda, Md., The Council, c1986. 382 p. (NCRP report, no. 86) QP82.2.E43N38 1986  
 "Issued April 2, 1986."  
 Bibliography: p. 290-353.

Nonionizing radiation protection. Edited by Michael J. Suess. Supported by the United Nations Environment Programme. Copenhagen, World Health Organization, Regional Office for Europe; Albany, N.Y., distributed by WHO Publications Centre USA, 1982. 267 p. QP82.2.R3N67 1982  
 Includes bibliographical references.  
 WHO regional publications. European series, no. 10.

Radiofrequency and microwaves. Geneva, World health Organization; Albany, N.Y., distributed by WHO Publications Centre USA, 1981. 134 p. (Environmental health criteria, 16) RA569.3.R3 1981  
 "Published under the joint sponsorship of the United Nations Environment Programme, the World Health Organization, and the International Radiation Protection Association."  
 Bibliography: p. 106-121.

Study in the USSR of medical effects of electric fields on electric power systems. Translations from Russian by G. G. Knickkrbocker. Presented as a public service by the Electrical Safety and Life Sciences Subcommittee of the Power and Environmental Sciences Committee of the IEEE Power Engineering Society. New York, Institute of Electrical and Electronics Engineers, c1975. 24 p. (Special publication of the IEEE Power Engineering Society, no. 10)

RC965.E4S78

### HANDBOOK

CRC handbook of biological effects of electromagnetic fields. Editors, Charles Polk and Elliot Postow. Boca Raton, Fla., CRC Press, c1986. 503 p. QP82.2.E43C73 1986\*  
 Includes bibliographies.

### BIBLIOGRAPHIES

B.C. Hydro. Library Services. Bibliography on biological effects of electric and magnetic fields, 1980-1984: literature held in B.C. Hydro Library. Vancouver, B.C., Library Services, B.C. Hydro, 1984. 16 p. Z6663.E44B18 1984

Current literature report on the carcinogenic properties of ionizing and non-ionizing radiation. Cincinnati, U.S. Dept. of Health, Education, and Welfare, Public Health Service, Center for Disease Control, National Institute for Occupational Safety and Health, Division of Biomedical and Behavioral Science; for sale by the Supt. of Docs., U.S. Govt. Print. Off., 1977-1978. 3 v. (DHEW (NIOSH) publication, no. 78-134) RC268.55.C86 and Pamphlet box (v. 2)\*  
 See particularly "Microwave and radiofrequency radiation": v. 2.

Heynick, Louise H. USAFAM review and analysis of radiofrequency radiation bioeffects literature. Second report. Menlo Park, Calif., SRI International, May 1982. 150 p. (SAM-TR-82-16) AD-A116 139\*\*  
 "Final report for period Sept. 1980-30 June 1981."

Heynick, Louise H., and John S. Krebs. USAFAM review and analysis of radiofrequency radiation bioeffects literature. Menlo Park, Calif., SRI International, Nov. 1981. 154 p. (SAM-TR-81-24) AD-A111 190\*\*  
 "Interim report for period 1 Mar. 1980-31 Aug. 1980."

\*\*Available in microform collection, Science Reading Room

Index of publications on biological effects of electromagnetic radiation (0-100 GHz.). Edited by James B. Kinn and Elliot Postow. Research Triangle, N.C., Health Effects Research Laboratory, U.S. Environmental Protection Agency, Mar. 1981. 575 p. PB81-181430\*\*  
"EPA-600/9-81-011."

Lockerby, Robert W. Biological effects of electromagnetic radiation. Monticello, Ill., Vance Bibliographies, 1981. 20 p. (Public administration series: bibliography, P-649) Z6663.E44L63 1981

Moore, Julie L. Cumulated index to the Bibliography of reported biological phenomena ('effects') and clinical manifestations attributed to microwave and radio-frequency radiation: report-supplements (no. 1-9), BEMS newsletter (B-1 through B-464), 1971-1981, compiled by Zorach R. Glaser. Indexed by Julie L. Moore. Riverside, Calif., J. Moore, 1984. 281 p. Z6663.M62M66 1984\*

See Appendix 1. "Chronology of the Bibliography and Supplements" to determine AD numbers for the Bibliography and Supplements.

See Appendix 2. "Listing of Bibliographic Supplements on 'Bioeffects of RF/Microwave Radiation' which have appeared in the Bioelectromagnetics Society's Newsletter." Pamphlet box\*

Reilly, J. Patrick. Human reactions to ELF electric and magnetic fields: an annotated bibliography of current literature. Laurel, Md., Johns Hopkins University, Applied Physics Laboratory, 1983. 88 p. QP82.2.E43R44 1983 ✓  
"JHU PPSE T-27."  
Prepared for the Maryland Power Plant Siting Program, July 1983.

----- Laurel, Md., Johns Hopkins University, Applied Physics Laboratory, 1984. 111 p. QP82.2.E43R44 1984  
"JHU PPSE T-30."  
Prepared for the Maryland Power Plant Siting Program, June 1984.

#### STATE-OF-ART-REVIEWS and CONFERENCE PROCEEDINGS

Biological effects of nonionizing radiation. Karl H. Illinger, editor. Washington, American Chemical Society, 1981. 342 p. (ACS symposium series, 157) QP82.2.R3B527  
"Based on a symposium sponsored by the Division of Physical Chemistry at the 179th meeting of the American Chemical Society, Houston, Texas, March 25-26, 1980."  
Includes bibliographical references.

Electrical and biological effects of transmission lines: a review. Prepared by the Biological Studies Task Team, Jack M. Lee, Jr. ... et al.. Portland, Ore., U.S. Dept. of Energy, Bonneville Power Administration, 1982. 93 p. QP82.2.E43E44 1982  
Bibliography: p. 79-93.  
"Second printing revised, Dec. 3, 1982."

Hanford Life Sciences Symposium, 18th, Richland, Wash., 1978. Biological effects of extremely low frequency electromagnetic fields: proceedings of the eighteenth annual Hanford Life Sciences Symposium at Richland, Washington, October 16-18, 1978. Sponsored by Office of Health and Environmental Research, U.S. Dept. of Energy, and Pacific North-West Laboratories, Battelle Memorial Institute, in cooperation with Electric Power Research Institute. Editors, Richard D. Phillips ... et al.. Oak Ridge, Tenn., Technical Information Center, U.S. Dept. of Energy, 1979. 577 p. (DOE symposium series, 50)

"CONF-781016."

QP82.2.E43H36 1978

Includes bibliographies.

Interaction between electromagnetic fields and cells. Edited by A. Chiabrera and C. Nicolini and H. P. Schwan. New York, Published in cooperation with NATO Scientific Affairs Division by Plenum Press, c1985. 640 p. (NATO Advanced Study Institute series. Series A, Life sciences, v. 97) QH656.I57 1985

"Proceedings of a NATO advanced research workshop entitled Interactions Between Electromagnetic Fields and Cells, held September 17-28, 1984 at the Ettore Majorana Centre for Scientific Culture in Erice, Sicily, Italy."

Includes bibliographies.

International School of Radiation Damage and Protection, 5th, Erice, Sicily, 1983. Biological effects and dosimetry of static and ELF electromagnetic fields. Edited by Martino Grandolfo, Sol M. Michaelson, and Alessandro Rindi. New York, Plenum Press, c1985. 697 p. (Ettore Majorana international science series. Life sciences, v. 19) QP82.2.E43I58 1983

"Proceedings of the fifth course of the International School of Radiation Damage and Protection of the Ettore Majorana Centre for Scientific Culture, held in November, 1983, in Erice, Sicily, Italy."

Includes bibliographies.

NATO Advanced Study Institute on Advances in Biological Effects and Dosimetry of Low Energy Electromagnetic Fields, Erice, Sicily, 1978. Biological effects and dosimetry of nonionizing radiation: radio-frequency and microwave energies. Edited by Martino Grandolfo, Sol M. Michaelson, and Alessandro Rindi. New York, Published in cooperation with NATO Scientific Affairs Division by Plenum Press, c1983. 669 p. (NATO Advanced Study Institute series. Series A, Life sciences, v. 49) QP82.2.M5N37 1981

"Proceedings of a NATO Advanced Study Institute on Advances in Biological Effects and Dosimetry of Low Energy Electromagnetic Fields which was also the fourth course of the International School of Radiation Damage and Protection of the Ettore Majorana Centre for Scientific Culture, held March 28-April 8, 1981, in Erice, Italy."

Includes bibliographical references.

Proceedings of a workshop on radiofrequency radiation bioeffects, 11-13 September 1984. Sponsored by Defense Research Group, Panel VIII, NATO AC/243. Editor, John C. Mitchell. Brooks Air Force Base, Tex., USAF School of Aerospace Medicine, Aerospace Medical Division (AFSC), April 1985. 233 p. (ULAFSAM-TP-85-14) AD-A157 090\*\*

"The workshop was held at the Research Establishment for Applied Science, Wachtberg-Werthoven, Federal Republic of Germany."

Symposium on Electromagnetic Fields in Biological Systems, Ottawa, Ont., 1978. Symposium on Electromagnetic Fields in Biological Systems: proceedings of a symposium held in Ottawa, Canada, June 27-30, 1978. Edited by Stanislaw S. Stuchly. Sponsored by the International Microwave Power Institute (IMPI) and the IEEE-Microwave Theory and Techniques Society, with the cooperation of the U.S. National Committee for the International Union of Radio Science and Canadian Commission A of URSI. Edmonton, Alta., International Microwave Power Institute, 1979. 323 p. Includes bibliographies. QP82.2.E43S95 1978

In many cases, conference proceedings or abstracts are published in journals. They can be identified through the abstracting and indexing publications listed in the ABSTRACTING AND INDEXING SERVICES section of this guide or through more specialized indexes to conference proceedings, such as Conference Papers Index (Z7403.C84\*) or Index to Scientific & Technical Proceedings (Z7403.I4\*). Some examples of conference material published in journals are given below.

Papers from the Symposium on Mechanisms of Bioelectrical Growth and Repair, appear in Journal of Bioelectricity, v. 3, no. 1/2, 1984: 3-101. QP341.J66

Translated abstracts of papers from the All-Union Symposium on the Biological Effects of Electromagnetic Fields, held in Puschino, U.S.S.R., in 1982, are contained in Biological Effects of Non-ionizing Electromagnetic Radiation, v. 9, Dec. 1984: 1-73. Pamphlet box\*

Proceedings of the First International Meeting of Association for the Biomedical Applications of Electromagnetism on Biological Effects and Therapeutic Applications of Electromagnetic Fields, held in Venice, Feb. 23-25, 1985, are contained in Bioelectrochemistry and Bioenergetics, v. 14, Nov. 1985: 3-244. QP341.B53

#### GOVERNMENT PUBLICATIONS

Biological effects of radiofrequency radiation. Edited by Joe A. Elder and Daniel F. Cahil. Research Triangle Park, N.C., Health Effects Research Laboratory, Office of Research and Development, U.S. Environmental Agency, 1984. 268 p. Pamphlet box\* and PB85-120848\*\*



Hankin, Norbert N. The radiofrequency radiation environment: environmental exposure levels and RF radiation emitting sources. Washington, Office of Radiation Programs, Office of Air and Radiation, U.S. Environmental Protection Agency, July 1986. 83 p. Pamphlet box\*  
Bibliography: p. 76-83.  
"EPA 520/1-85-014."

United States. Congress. House. Committee on Science and Technology. Subcommittee on Natural Resources and Environment. Research on health effects of nonionizing radiation: hearing ... Ninety-sixth Congress, first session, July 12, 1979. Washington, U.S. Govt. Print. Off., 1979. 457 p. KF27.S398 1979g  
"No. 70."  
Includes bibliographical references.

United States. Congress. House. Committee on Science and Technology. Subcommittee on Investigations and Oversight. Potential health effects of video display terminals and radio frequency heaters and sealers: hearings ... Ninety-seventh Congress, first session, May 12, 13, 1981. Washington, U.S. Govt. Print. Off., 1981. 764 p. KF27.S3975 1981c  
"No. 13."  
Bibliography: p. 682-686.

United States. Congress. Senate. Committee on Commerce, Science, and Transportation. Microwave irradiation of the U.S. Embassy in Moscow: review of its history and studies to determine whether or not related health defects were experienced by employees assigned in the period, 1953-1977. Prepared at the request of Howard W. Cannon, chairman, Committee on Commerce, Science, and Transportation, United States Senate. Washington, U.S. Govt. Print. Off., 1979. 26 p. JX1706.A58R88  
"At head of title: 96th Congress, 1st session. Committee print."

ABSTRACTING AND INDEXING SERVICES that index relevant journal articles and other literature are listed below. Some terms are suggested as aids in searching.

Abstracts on Health Effects of Environmental Pollutants (1972-)  
RA565.A1A26\*

See: Keyword of interest, e.g., Electromagnetic (Field(s), Frequency, etc.), Microwave (Exposure, Radiation, etc.)

Applied Science & Technology Index (1913-) 27913.17\*

See: Electromagnetic Fields--Physiological Effect  
Microwaves--Physiological Effect  
Radio Waves--Physiological Effect

Biological Abstracts (1926-) QH301.B37\* and Biological Abstracts/RRM (1967-) Z53211.B672\*

See: Keyword of interest, e.g. Electromagnetic (Field(s) Exposure, Radiation, etc.), Extremely-Low-Frequency, ELF, Nonionizing Radiation, Radiofrequency (Exposure, Radiation, etc.), RF

Biological & Agricultural Index (1916-) Z5073.A46\*

See: Electromagnetic Waves--Biological Effect  
Microwaves--Biological Effect

Environmental Periodicals Bibliography (1972-) Z5863.E57E58\*

See: Heading beginning Electromagnetic; Microwave; Radiation,  
Nonionizing; Radiofrequency

Index Medicus (1960-) Z6660.I422\*

See: Electromagnetics  
Microwaves  
Radiation, Non-ionizing  
Radio Waves

INIS Atomindex (1969-) Z7144.N8I18\*

See: Electromagnetic Radiation--Biological Radiation effects  
Electromagnetic Radiation--Dose Limits  
Electromagnetic Radiation--Radiation Protection  
Long Wave Radiation  
Microwave Radiation--Biological Effects  
Microwave Radiation--Biological Radiation Effects  
Microwave Radiation--Prenatal Irradiation  
Microwave Radiation--Pulsed Irradiation  
Microwave Radiation--Radiation Protection

Magazine Index 1976-)

See: Electromagnetic Fields  
Electromagnetic Waves--Environmental Aspects  
Electromagnetic Waves--Medical Use  
Electromagnetic Waves--Physiological Aspects  
Microwaves--Environmental Aspects  
Microwaves--Medical Use  
Microwaves--Physiological Aspects  
Radio Waves

Readers' Guide to Periodical Literature (1900-) AI3.R45

See: Electromagnetic Fields--Physiological Effects  
Electromagnetic Therapy  
Electromagnetic Waves--Physiological Effects  
Microwaves--Physiological Effects

JOURNALS that often contain articles on physiological and health effects  
of electromagnetic radiation are

Bioelectromagnetics QP82.2.E43B53

Bioelectrochemistry and bioenergetics QP341.B53

Journal of bioelectricity QP341.J66

Microwave news (current issues only in LM133)

USSR Report. Life Sciences. Effects of Nonionizing Electromagnetic  
Radiation (JPRS) (available in LM133 and LJ140A)

VDT news (review before binding in LM133)

## REPRESENTATIVE JOURNAL ARTICLES

- Adey, W. R. Tissue interactions with nonionizing electromagnetic fields. *Physiological review*, v. 61, 1981: 435-514. QP1.P45
- Bauchinger, M., and others. Analysis of structural chromosome changes and SCE after occupational long-term exposure to electric and magnetic fields from 380 kV-systems. *Radiation and environmental biophysics*, v. 19, 1981: 235-238. QH505.A1B57
- Blackman, C. F. Effects of ELF (1-120 Hz) and modulated (50 Hz) RF fields on the efflux of calcium ions from brain tissue in vitro. *Bioelectromagnetics*, v. 6, 1985: 1-11. QP82.2.E43B53
- Delgado, J. M. R., and others. Embryological changes induced by weak, extremely low frequency electromagnetic fields. *Journal of anatomy*, v. 134, 1982: 533-551. QL801.J7
- Justesen, Don R. Behavioral and psychological effects of microwave radiation. *In* New York Academy of Medicine. *Bulletin*, v. 55, 1980: 1058-1078. R15.N62
- Lerner, Eric J. The drive to regulate electromagnetic fields. *IEEE spectrum*, 21, Mar. 1984: 63-70. TK1.I15 and Pamphlet box\*
- Nordstrom, S., E. Birke, and L. Gustavson. Reproductive hazards among workers at high voltage substations. *Bioelectromagnetics*, v. 4, 1983: 91-101. QP82.2.E43B53
- Wertheimer, Nancy, and Ed Leeper. Possible effects of electric blankets and heated waterbeds on fetal development. *Bioelectromagnetics*, v. 7, 1986: 13-22. QP82.2.E43B53

REPORTS and other types of literature are indexed in the following guides:

Government Reports Announcements & Index (1946-) Z7916.G78\*

See: Electromagnetic Fields  
Microwaves  
Radiation Protection  
Radiofrequency

Monthly Catalog of United States Government Publications (1895-)  
Z1223.A18\*

See: Electromagnetism--Physiological Effect  
Microwaves--Physiological Effect  
Microwaves--Psychological Aspects  
Radiation--Physiological Effect

**SELECTED TECHNICAL REPORTS**, sold by the National Technical Information Service, Springfield, Virginia 22161, include the following:

- Assessments and viewpoints on the biological and human health effects of extremely low frequency (ELF) electromagnetic fields. Arlington, Va., American Institute of Biological Sciences, May 1985. 461 p.  
AD-A156 942\*\*
- Behavioral effects of microwave radiation absorption. Editors, John C. Monahan and John A. D'Andrea. Rockville, Md., Center for Devices and Radiological Health, Food and Drug Administration, Aug. 1985. 173 p. (FDA85-8238)  
PB86-113735\*\*
- Biological and human health effects of extremely low frequency electromagnetic fields: post-1977 literature review. Arlington, Va., American Institute of Biological Sciences, Mar. 1985. 327 p.  
AD-A152 731\*\*
- Blackman, Carl F. Biological influences of low-frequency sinusoidal electromagnetic signals alone and superimposed on RF carrier waves. Research Triangle Park, N.C., Health Effects Research Laboratory, U.S. Environmental Protection Agency, Jan. 1985. 15 p.  
(EPA/600/D-85-010)  
PB85-161248\*\*
- Meltz, Martin L., and Kathleen A. Walker. Genetic effects of microwave exposure on mammalian cells in vitro: volume 1. San Antonio, Tex., University of Texas Health Science Center, June 1984. 39 p.  
(USAFSAM-TR-84-22)  
AD-A145 946\*\*
- Meltz, Martin L., Clifton R. Harris, and Kathleen A. Walker. Genetic effects of microwave exposure on mammalian cells in vitro: volume 2. San Antonio, Tex., University of Texas Health Science Center, Oct. 1984. 88 p. (USAFSAM-TR-84-24)  
AD-A149 347\*\*
- O'Connor, Mary Ellen, and Robert Strattan. Teratogenic effects of microwave radiation. Tulsa, Okl., Departments of Psychology and Electrical Engineering, University of Tulsa, May 1985. 42 p.  
(EPA/600/1-85/007)  
PB85-207462\*\*
- Raines, Jeremy K. Electromagnetic field interaction with the human body: observed facts and theories. Greenbelt, Md., Goddard Space Flight Center, National Aeronautics and Space Administration, Apr. 1981. 123 p. (NASA-CR-166661)  
N81-25668\*\*

**SELECTED MATERIALS** available in the Science Reading Room pamphlet boxes include:

- Bibliography of reported biological, phenomena ('effects') and clinical manifestations attributed to microwave and radio-frequency radiation. Compiled by Zorach R. Glaser. Bethesda, Md., Naval Medical Research Institute, National Naval Medical Center, 1971. 92 p.

Dodge, Christopher H. Video display terminals: the controversy about health and safety issues. Washington, Congressional Research Service, Library of Congress, 1984 (updated Jan. 30, 1985). 20 p.

Environmental Protection Agency. Federal radiation protection guidance: proposed alternatives for controlling public exposure to radiofrequency radiation; notice of proposed recommendations. Federal register, July 30, 1986: 27318-27339.

Gannon, Robert. Electromagnetic pollution: are they zapping you? Popular science, v. 223, Dec. 1983: 98-101, 176.

The health hazards of computers: a guide to worrying intelligently. Whole earth review, no. 48, Fall 1985: 80-93.

Lerner, Eric J. The drive to regulate electromagnetic fields. IEEE spectrum, v. 21, Mar. 1984: 63-70.

McAuliffe, Kathleen. The mind fields. Omni, v. 7, Feb. 1985: 41-42, 44, 96, 98, 100-104.

Miller, Morton W. and Gary E. Kaufman. High voltage overhead. Environment, v. 20, Jan./Feb. 1978: 6-15, 32-36.

Morgan, Granger M. Risk research: when should we say "enough"? Science, v. 232, May 23, 1986: 917.

Nichols, Christina. Power towers: are high-voltage lines dangerous? Environmental action, v. 17, May/June 1986: 26-27.

#### ADDITIONAL SOURCES OF INFORMATION

Center for Devices and Radiological Health  
U.S. Food and Drug Administration  
5600 Fishers Lane  
Rockville, Maryland 20857  
Telephone: (301) 443-4690

National Council on Radiation Protection and Measurements (NCRP)  
7910 Woodmont Ave., Suite 1016  
Bethesda, Maryland 20814  
Telephone: (301) 657-2652

Office of Radiation Programs  
Office of Air and Radiation  
U.S. Environmental Protection Agency  
401 M. Street, S. W.  
Washington, D.C. 20460  
Telephone: (202) 475-9605

Office of Spectrum Management  
National Telecommunications and Information Administration  
14th Street and Constitution Avenue, N.W.  
Washington, D.C. 20230  
Telephone: (202) 377-1850

Office of Radiation Programs Technical Publications  
Nonionizing Radiation

Publications of the Office of Radiation Programs are available from the National Technical Information Service (NTIS), Springfield, VA 22161. Current prices should be obtained directly from NTIS using the indicated NTIS Order number. Single copies of some of the publications listed below may also be available without charge from the Office of Radiation Programs (AW-461), 401 M St., SW Washington, DC 20460.

EPA ORP/SID 72-3	Reference Data for Radiofrequency Emission Hazard Analysis (NTIS Order No. PB 220 471)
EPA/ORP 73-2	Environmental Exposure to Nonionizing Radiation, (Available NTIS only, Order No. PB 220 851)
EPA-520/2-73-001	Nonionizing Measurement Capabilities: State and Federal Agencies (Available NTIS only, Order No. PB 226 778/AS)
EPA-520/1-74-005	RF Pulse Spectral Measurements in the Vicinity of Several ATC Radars (NTIS Order No. PB 235 733)
EPA-520/2-74-008	An Evaluation of Satellite Communication Systems as Sources of Environmental Microwave Radiation (NTIS Order No. PB 257 138/AS)
ORP/EAD 75-1	An Analysis of Broadcast Radiation Levels in Hawaii (NTIS Order No. PB 261 316/AS)
ORP/EAD-76-1	Radiation Characteristics of Traffic Radar Systems (NTIS Order No. PB 257 077/AS)
ORP/EAD-76-2	A Measurement of RF Field Intensities in the Immediate Vicinity of an FM Broadcast Station Antenna (NTIS Order No. PB 257 698/AS)
EPA-520/2-76-008	An Examination of Electric Fields Under EHV Overhead Power Transmission Lines (NTIS Order No. PB 270 613/AS)
ORP/EAD-77-2	An Investigation of Broadcast Radiation Intensities at Mt. Wilson, California (NTIS Order No. PB 275 040/AS)
ORP/EAD-77-3	An Analysis of Radar Exposure in the San Francisco Area (NTIS Order No. PB 273 188/AS)